

# involve

a journal of mathematics

## Existence of positive solutions for an approximation of stationary mean-field games

Nojood Almayouf, Elena Bachini, Andreia Chapouto, Rita Ferreira,  
Diogo Gomes, Daniela Jordão, David Evangelista Junior,  
Avetik Karagulyan, Juan Monasterio, Levon Nurbekyan,  
Giorgia Pagliar, Marco Piccirilli, Sagar Pratapsi, Mariana Prazeres,  
João Reis, André Rodrigues, Orlando Romero, Maria Sargsyan,  
Tommaso Seneci, Chuliang Song, Kengo Terai, Ryota Tomisaki,  
Hector Velasco-Perez, Vardan Voskanyan and Xianjin Yang



# Existence of positive solutions for an approximation of stationary mean-field games

Nojood Almayouf, Elena Bachini, Andreia Chapouto, Rita Ferreira,  
Diogo Gomes, Daniela Jordão, David Evangelista Junior,  
Avetik Karagulyan, Juan Monasterio, Levon Nurbekyan,  
Giorgia Pagliar, Marco Piccirilli, Sagar Pratapsi, Mariana Prazeres,  
João Reis, André Rodrigues, Orlando Romero, Maria Sargsyan,  
Tommaso Seneci, Chuliang Song, Kengo Terai, Ryota Tomisaki,  
Hector Velasco-Perez, Vardan Voskanyan and Xianjin Yang

(Communicated by Kenneth S. Berenhaut)

Here, we consider a regularized mean-field game model that features a low-order regularization. We prove the existence of solutions with positive density. To do so, we combine a priori estimates with the continuation method. In contrast with high-order regularizations, the low-order regularizations are easier to implement numerically. Moreover, our methods give a theoretical foundation for this approach.

## 1. Prologue

On August 22, 2015, eighteen young mathematicians (B.Sc. and M.Sc. students) arrived at King Abdullah University of Science and Technology (KAUST) in Thuwal, Kingdom of Saudi Arabia. They were participants in the first KAUST summer camp in applied partial differential equations. Among them were Argentinians, Armenians, Chinese, Italians, Japanese, Mexicans, Portuguese, and Saudis. For many of them, this was their first time abroad. All were looking forward to the following three weeks.

We designed the summer camp to give an intense hands-on three-week Ph.D. experience. It comprised courses, seminars, a project, and a final presentation. The

---

*MSC2010:* 49L25, 91A13, 35J87.

*Keywords:* mean-field games, low-order regularizations, monotone methods, positive solutions.

Rita Ferreira, Diogo Gomes, David Evangelista Junior, Levon Nurbekyan, Mariana Prazeres, Vardan Voskanyan, and Xianjin Yang were partially supported by KAUST baseline and start-up funds and KAUST SRI, Uncertainty Quantification Center in Computational Science and Engineering. The other authors were partially supported by KAUST Visiting Students Research Program.

project was an essential component of the summer camp, and its main outcome is the present paper. Our objectives were to introduce students to an active research topic, teach effective paper writing techniques, and develop their presentation skills. Numerous challenges lay ahead. First, we had three weeks to achieve these goals. Second, students had distinct backgrounds. Third, we planned to study a research-level problem, not a simple exercise.

We selected a problem in mean-field games, a recent and active area of research. The primary goal was to prove the existence of solutions of a system of partial differential equations. To avoid unnecessary technicalities, we considered the one-dimensional case, where the partial differential equations become ordinary differential equations. The project involved partial differential equation methods that are usually taught in advanced courses: a priori estimate methods, the infinite-dimensional implicit function theorem, and the continuation method. In spite of the elementary nature of the proofs, the results presented here are a relevant and original contribution to the theory of mean-field games.

We divided the students into five groups and assigned tasks to each of them. Roughly, each of the sections of this paper corresponds to a task. The students were given a rough statement of the results to be proven, and their task was to figure out the appropriate assumptions, the precise statements, and the proofs. The work of the different groups had to be coordinated to make sure that the assumptions, results, and proofs fit nicely with each other and that duplicate work was avoided. Several KAUST graduate students and postdocs were of invaluable help in this regard.

This project would not have been possible within such a short time frame without the use of new technologies. The paper was written in a collaborative fashion using the platform [Authorea](#) that allowed all the groups to work simultaneously. In this way, all groups had access to the latest version of the assumptions and to the current statements of the theorems and propositions. Each group could easily comment and make corrections on other group's work.

This project illustrates how research in mathematics can be a collaborative experience even with a large number of participants. Moreover, it gave each of the students in the summer camp a glimpse of real research in mathematics. Finally, this was the first experience for the Ph.D. students and postdocs who helped in this project in mentoring and advising students. This summer camp was a unique and valuable experience for all participants whose results we share in this paper.

## 2. Introduction

Mean-field game (MFG) theory is the study of strategic decision making in large populations of small interacting individuals who are also called agents or players. The MFG framework was developed in the engineering community by Caines,

Huang, and Malhamé [Huang et al. 2006; 2007] and in the mathematical community by Lasry and Lions [2006a; 2006b; 2007]. These games model the behavior of rational agents who play symmetric differential games. In these problems, each player chooses their optimal strategy in view of global (or macroscopic) statistical information on the ensemble of players. This approach leads to novel problems in nonlinear equations. Current research topics are the applications of MFGs (including, for example, growth theory in economics and environmental policy), mathematical problems related to MFGs (existence, uniqueness, and regularity questions), and numerical methods in the MFGs framework (discretization, convergence, and efficient implementation).

Here, we consider the following problem:

**Problem 1.** Let  $\mathbb{T} = \mathbb{R}/\mathbb{Z}$  denote the one-dimensional torus, identified with the interval  $[0, 1]$  whenever convenient. Fix a  $C^2$  Hamiltonian,  $H : \mathbb{R} \rightarrow \mathbb{R}$ , and a continuous potential,  $V : \mathbb{T} \rightarrow \mathbb{R}$ . Let  $\alpha$  and  $\epsilon$  be positive numbers with  $\epsilon \leq 1$  for definedness. Find  $u, m \in C^2(\mathbb{T})$  satisfying  $m > 0$  and

$$\begin{cases} u - u_{xx} + H(u_x) + V(x) = m^\alpha + \epsilon(m - m_{xx}), \\ m - m_{xx} - (H'(u_x)m)_x = 1 - \epsilon(u - u_{xx}). \end{cases} \quad (2-1)$$

In this problem,  $m$  is the distribution of players and  $u(x)$  is the value function for a typical player in the state  $x$ . We stress that the condition  $m > 0$  is an essential component of the problem. So, if  $(u, m)$  solves **Problem 1**, we require  $m$  to be strictly positive. We will show the existence of solutions to this problem under suitable assumptions on the Hamiltonian that are described in **Section 3**. An example that satisfies those assumptions is  $H(p) = (1 + p^2)^{\gamma/2}$  with  $1 < \gamma < 2$  and any  $V : \mathbb{T} \rightarrow \mathbb{R}$  of class  $C^2$ .

When  $\epsilon = 0$ , (2-1) becomes

$$\begin{cases} u - u_{xx} + H(u_x) + V(x) = m^\alpha, \\ m - m_{xx} - (H'(u_x)m)_x = 1. \end{cases} \quad (2-2)$$

The system in (2-2) is a typical MFG model similar to the one introduced in [Lasry and Lions 2006a]. The Legendre transform of the Hamiltonian,  $H$ , given by  $L(v) = \sup_p -pv - H(p)$  is the cost in units of time that an agent incurs by choosing to move with a drift  $v$ ; the potential  $V$  accounts for spatial preferences of the agents; the term  $m^\alpha$  encodes congestion effects.

The MFG models proposed in [Lasry and Lions 2006a; 2006b] consist of a system of partial differential equations that have (2-2) as a particular case. The current literature covers a broad range of problems, including stationary problems [Gomes et al. 2012; 2014; Gomes and Ribeiro 2013; Gomes and Sánchez Morgado 2014; Pimentel and Voskanyan 2015], heterogeneous populations [Cirant 2015], time-dependent models [Cardaliaguet et al. 2015; Gomes et al. 2015; 2016; Gomes

and Pimentel 2015; 2016; Porretta 2014; 2015], congestion problems [Gomes and Mitake 2015; Graber 2015], and obstacle-type problems [Gomes and Patrizi 2015]. For a recent account of the theory of MFGs, we suggest the survey paper [Gomes and Saúde 2014] and the course [Lions 2012].

The system in (2-1) arises as an approximation of (2-2) that preserves monotonicity properties. Monotonicity-preserving approximations to MFG systems were introduced in [Ferreira and Gomes 2015]. In that paper, the authors consider mean-field games in dimension  $d \geq 1$ , which include the following example:

$$\begin{cases} u - \Delta u + H(Du, x) + V(x) = m^\alpha + \epsilon(m + \Delta^{2q}m) + \beta_\epsilon(m), \\ m - \Delta m - \operatorname{div}(D_p H(Du, x)m) = 1 - \epsilon(u + \Delta^{2q}u), \end{cases} \quad (2-3)$$

where  $q$  is a large enough integer, and  $\beta_\epsilon$  is a suitable penalization that satisfies  $\beta_\epsilon(m) \rightarrow -\infty$  as  $m \rightarrow 0$ . Then, as  $\epsilon \rightarrow 0$ , the solutions of (2-3) converge to solutions of (2-2). Yet, from the perspective of numerical methods, both the high-order degree of (2-3) and the singularity caused by the penalty,  $\beta_\epsilon$ , are unsatisfactory due to a poor conditioning of discretizations. Here, we investigate a low-order regularization that may be more suitable for computational problems.

A fundamental difficulty in the analysis of (2-1) is the nonnegativity of  $m$ . The Fokker–Planck equation in (2-2) has a maximum principle, and, consequently,  $m \geq 0$  for any solution of (2-2). Due to the coupling, this property is not evident in the corresponding equation in (2-1). The previous regularization in (2-3) relies on a penalty that forces the positivity of  $m$ . This mechanism does not exist in (2-1), and we are not aware of any general method to prove the existence of positive solutions of (2-1).

Our main result is the following theorem.

**Theorem 2.1.** *Suppose that Assumptions 1–7 hold (see Section 3). Then, there exists  $\epsilon_0 > 0$  such that for all  $0 < \epsilon < \epsilon_0$ , Problem 1 admits a  $C^{2,1/2}$  solution  $(u, m)$ .*

Theorem 2.1 introduces a low-order regularization procedure for (2-2) for which existence of solutions can be established without penalty terms. Because high-order regularization methods and penalty terms create serious difficulties in the numerical implementation, this result is relevant to the numerical approximation of (2-2). Moreover, we believe that the techniques we consider here can be extended to higher-dimensional problems.

To prove the main result, we use the continuation method. The first step is to establish a priori estimates for the solutions of (2-1). Then, we replace the potential  $V$  by  $\lambda V$  for  $0 \leq \lambda \leq 1$ . For  $\lambda = 0$ , which corresponds to  $V = 0$  in (2-1), we determine an explicit solution. The a priori estimates give that the set  $\Lambda$  of values  $\lambda$  for which (2-1) has a solution is a closed set. Finally, we apply an infinite-dimensional version of the implicit function theorem to show that  $\Lambda$  is relatively open in  $[0, 1]$ . This proves the existence of solutions.

The remainder of this paper is structured as follows. We discuss the main assumptions in [Section 3](#). Next, in [Section 4](#), we start our study of (2-1) by considering the case  $V = 0$  and constructing an explicit solution. Sections 5–9 are devoted to a priori estimates for solutions of (2-1). These estimates include energy and second-order bounds, discussed respectively in Sections 5 and 6, Hölder and  $C^{2,1/2}$  estimates, addressed respectively in Sections 7 and 8, and lower bounds on  $m$ , given in [Section 9](#). Next, we lay out the main results needed for the implicit function theorem. We introduce the linearized operator in [Section 10](#) and discuss its injectivity and surjectivity properties. Finally, the proof of [Theorem 2.1](#) is presented in [Section 11](#).

### 3. Main assumptions

We start by recalling that  $C^{2,1/2}(\mathbb{T})$  is the space of all functions in  $C^2(\mathbb{T})$  whose second derivative is  $\frac{1}{2}$ -Hölder continuous.

To prove [Theorem 2.1](#), we need to introduce various assumptions that are natural in this class of problems. These encode distinct properties of the Hamiltonian in a convenient way. We begin by stating a polynomial growth condition for the Hamiltonian.

**Assumption 1.** There exist positive constants,  $C_1, C_2, C_3$ , and  $\gamma > 1$ , such that for all  $p \in \mathbb{R}$ , the Hamiltonian  $H$  satisfies

$$-C_1 + C_2|p|^\gamma \leq H(p) \leq C_1 + C_3|p|^\gamma.$$

For convex Hamiltonians, the expression  $pH'(p) - H(p)$  is the Lagrangian written in momentum coordinates. The next assumption imposes polynomial growth in this quantity.

**Assumption 2.** There exist positive constants,  $\tilde{C}_1, \tilde{C}_2$ , and  $\tilde{C}_3$ , such that for all  $p \in \mathbb{R}$ , we have

$$-\tilde{C}_1 + \tilde{C}_2|p|^\gamma \leq pH'(p) - H(p) \leq \tilde{C}_1 + \tilde{C}_3|p|^\gamma.$$

Because we look for solutions  $(u, m) \in C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T})$  of [Problem 1](#), we require in Assumptions 3 and 5 more regularity for  $V$  and  $H$ .

**Assumption 3.** The potential  $V$  is of class  $C^2$ .

Because the Hamilton–Jacobi equation in (2-2) arises from an optimal control problem, it is natural to suppose that the Hamiltonian  $H$  is convex.

**Assumption 4.**  $H$  is convex.

**Assumption 5.** The Hamiltonian  $H$  is of class  $C^4$ .

Here, we work with subquadratic Hamiltonians. Accordingly, we impose the following condition on  $\gamma$ .

**Assumption 6.** The constant  $\gamma$  satisfies  $\gamma < 2$ .

Finally, we state a growth condition on the derivative of the Hamiltonian. The exponent  $\gamma$  is the same as in Assumptions 1 and 2. This is a natural growth condition that the model  $H(p) = (1 + |p|^2)^{\gamma/2}$  satisfies.

**Assumption 7.** There exists a positive constant,  $\bar{C}$ , such that for all  $p \in \mathbb{R}$ , we have

$$|H'(p)| \leq \bar{C}(1 + |p|^{\gamma-1}).$$

#### 4. The $V = 0$ case

To prove Theorem 2.1, we use the continuation method. More precisely, we consider system (2-1) with  $V$  replaced by  $\lambda V$  for  $0 \leq \lambda \leq 1$ . Next, we show the existence of the solution for all  $0 \leq \lambda \leq 1$ . As a starting point, we study the  $\lambda = 0$  case; that is,  $V = 0$ . We show that (2-1) admits a solution in this particular instance.

**Proposition 4.1.** *Suppose that  $V = 0$ . Then, there exists an  $\epsilon_0 > 0$  such that for all  $0 < \epsilon < \epsilon_0$ , Problem 1 admits a solution  $(u, m)$ .*

*Proof.* We look for constant solutions  $(u, m)$ . In this case, we have  $u_x = u_{xx} = m_x = m_{xx} = 0$ . Accordingly, (2-1) reduces to

$$\begin{cases} u + H(0) = m^\alpha + \epsilon m, \\ m = 1 - \epsilon u. \end{cases}$$

In the previous system, solving the first equation for  $u$  and replacing the resulting expression into the second, we get

$$\epsilon m^\alpha + (1 + \epsilon^2)m - 1 - \epsilon H(0) = 0. \tag{4-1}$$

We set  $g(m) = \epsilon m^\alpha + (1 + \epsilon^2)m - 1 - \epsilon H(0)$ , so that (4-1) reads  $g(m) = 0$ . Next, we notice that  $g(0) = -1 - \epsilon H(0)$ . For small enough  $\epsilon_0 > 0$  and for all  $0 < \epsilon < \epsilon_0$ , we have  $g(0) < 0$ . On the other hand, if we take a constant  $C > |H(0)|$ , we have

$$g(1 + \epsilon C) > 1 + \epsilon C - 1 - \epsilon H(0) = \epsilon(C - H(0)) > 0.$$

Because  $0 < 1 + \epsilon C$ , by the intermediate value theorem, there exists a constant  $m_0 \in ]0, 1 + \epsilon C[$  such that  $g(m_0) = 0$ . Then, setting  $u_0 = (1 - m_0)/\epsilon$ , we conclude that the pair  $(u_0, m_0)$  satisfies the requirements. □

**Remark 4.2.** Note that if  $H(0) > 0$ , then  $g(0) < 0$  and  $g(1 + \epsilon C) > 0$ . In this case, the previous proposition holds for all  $\epsilon > 0$ .

## 5. Energy estimates

MFG systems such as (2-2) admit many a priori estimates. Among those, energy estimates stand out for their elementary proof—the multiplier method. Here, we apply this method to (2-1).

**Proposition 5.1.** *Suppose that Assumptions 1 and 2 hold. Let  $(u, m)$  solve Problem 1. Then,*

$$\int_0^1 m^{\alpha+1} dx + \int_0^1 |u_x|^\gamma (1+m) dx + \epsilon \int_0^1 (u^2 + m^2 + u_x^2 + m_x^2) dx \leq C, \quad (5-1)$$

where  $C$  is a universal positive constant depending only on the constants in Assumptions 1 and 2 and on  $\|V\|_{L^\infty}$ .

*Proof.* We begin by multiplying the first equation in (2-1) by  $(1 + \epsilon - m)$  and the second one by  $u$ . Adding the resulting expressions and integrating, we get

$$\begin{aligned} & \int_0^1 [(1+\epsilon)H(u_x) + m(u_x H'(u_x) - H(u_x))] dx \\ & \quad + \int_0^1 m^{\alpha+1} dx + \epsilon \int_0^1 (u^2 + m^2 + u_x^2 + m_x^2) dx \\ = & -\epsilon \int_0^1 u dx + \int_0^1 (m-1-\epsilon)V(x) dx + (1+\epsilon) \int_0^1 m^\alpha dx + \epsilon(1+\epsilon) \int_0^1 m dx, \end{aligned} \quad (5-2)$$

where we used integration by parts and the periodicity of  $u$  and  $m$  to obtain

$$\begin{aligned} & \int_0^1 m u_{xx} dx - \int_0^1 u m_{xx} dx = 0, \\ & \int_0^1 u_{xx} dx = u_x|_0^1 = 0, \quad \int_0^1 m_{xx} dx = m_x|_0^1 = 0, \\ & \int_0^1 m m_{xx} dx = - \int_0^1 m_x^2 dx, \quad \int_0^1 u u_{xx} dx = - \int_0^1 u_x^2 dx, \\ & \int_0^1 u (H'(u_x) m)_x dx = - \int_0^1 u_x H'(u_x) m dx. \end{aligned}$$

Next, we observe that by Assumptions 1 and 2, and using the fact that  $0 < \epsilon \leq 1$ , we have

$$\begin{aligned} & \int_0^1 [(1+\epsilon)H(u_x) + m(H'(u_x)u_x - H(u_x))] dx \\ & \quad \geq \int_0^1 [-2C_1 - \tilde{C}_1 m + K_0 |u_x|^\gamma (1+m)] dx, \end{aligned} \quad (5-3)$$

where  $K_0 := \min\{C_2, \tilde{C}_2\}$ .



From (5-2) and (5-3), it follows that

$$\begin{aligned} & \int_0^1 K_0|u_x|^\gamma(1+m) \, dx + \int_0^1 m^{\alpha+1} \, dx + \epsilon \int_0^1 (u^2+m^2+u_x^2+m_x^2) \, dx \\ & \leq \frac{\epsilon}{2} \int_0^1 u^2 \, dx + \frac{1}{2} + (\|V\|_\infty + 2 + \tilde{C}_1) \int_0^1 m \, dx + 2 \int_0^1 m^\alpha \, dx + 2(\|V\|_\infty + C_1), \end{aligned} \tag{5-4}$$

where we also used the estimates  $2u \leq u^2 + 1$  and  $0 < \epsilon \leq 1$ .

Finally, we observe that for every  $\delta_1, \delta_2 > 0$ , there exist constants,  $K_1$  and  $K_2$ , such that

$$\int_0^1 m^\alpha \, dx \leq \delta_1 \int_0^1 m^{\alpha+1} \, dx + K_1, \quad \int_0^1 m \, dx \leq \delta_2 \int_0^1 m^{\alpha+1} \, dx + K_2. \tag{5-5}$$

Consequently, taking  $\delta_1 = \frac{1}{8}$  and  $\delta_2 = 1/(4(\|V\|_\infty + 2 + \tilde{C}_1))$  in (5-5) and using the resulting estimates in (5-4), we conclude that (5-1) holds.  $\square$

**Corollary 5.2.** *Suppose that Assumptions 1 and 2 hold. Let  $(u, m)$  solve Problem 1. Then,*

$$\int_0^1 m \, dx \leq C,$$

where  $C$  is a universal positive constant depending only on the constants in Assumptions 1 and 2 and on  $\|V\|_{L^\infty}$ .

*Proof.* Due to (5-1) and because  $m$  is positive,

$$\int_0^1 m^{\alpha+1} \, dx \leq C,$$

where  $C$  is a universal positive constant depending only on the constants in Assumptions 1 and 2 and on  $\|V\|_{L^\infty}$ . Consequently, using Young’s inequality, we have

$$\int_0^1 m \, dx \leq \frac{1}{\alpha+1} \int_0^1 m^{\alpha+1} \, dx + \frac{\alpha}{\alpha+1} \leq \frac{C}{\alpha+1} + \frac{\alpha}{\alpha+1}. \quad \square$$

### 6. Second-order estimates

We proceed in our study of (2-1) by examining another technique to obtain a priori estimates. These estimates give additional control over high-order norms of the solutions.

**Proposition 6.1.** *Suppose that Assumption 3 holds. Let  $(u, m)$  solve Problem 1. Then, we have*

$$\int_0^1 (H''(u_x)u_{xx}^2m + \alpha m^{\alpha-1}m_x^2) \, dx + \epsilon \int_0^1 (m_x^2 + m_{xx}^2 + u_x^2 + u_{xx}^2) \, dx \leq C, \tag{6-1}$$

where  $C > 0$  denotes a universal constant depending only on  $\|V\|_{C^2}$ . Moreover, under Assumption 4,

$$\int_0^1 \alpha m^{\alpha-1} m_x^2 \, dx + \epsilon \int_0^1 (m_x^2 + m_{xx}^2 + u_x^2 + u_{xx}^2) \, dx \leq C. \tag{6-2}$$

*Proof.* To simplify the notation, we represent by  $C$  any positive constant that depends only on  $\|V\|_{C^2}$  and whose value may change from one instance to another.

Multiplying the first equation in (2-1) by  $m_{xx}$  and the second one by  $u_{xx}$  yields

$$\begin{aligned} (u - u_{xx} + H(u_x) + V(x))m_{xx} &= (m^\alpha + \epsilon(m - m_{xx}))m_{xx}, \\ (m - m_{xx} - (H'(u_x)m)_x)u_{xx} &= (1 - \epsilon(u - u_{xx}))u_{xx}. \end{aligned}$$

Subtracting the above equations integrated over  $[0, 1]$  gives

$$\begin{aligned} &\int_0^1 (um_{xx} - mu_{xx} + u_{xx}) \, dx + \int_0^1 [H(u_x)m_{xx} + (H'(u_x)m)_x u_{xx}] \, dx \\ &+ \int_0^1 V(x)m_{xx} \, dx - \int_0^1 m^\alpha m_{xx} \, dx + \epsilon \int_0^1 (-mm_{xx} + m_{xx}^2 - uu_{xx} + u_{xx}^2) \, dx = 0. \end{aligned} \tag{6-3}$$

Next, we evaluate each of the integrals above. Using the integration by parts formula and the periodicity of boundary conditions, we have

$$\int_0^1 (um_{xx} - mu_{xx} + u_{xx}) \, dx = 0. \tag{6-4}$$

In addition,

$$\begin{aligned} &\int_0^1 [(H'(u_x)m)_x u_{xx} + H(u_x)m_{xx}] \, dx \\ &= \int_0^1 [H''(u_x)mu_{xx}^2 + (H(u_x))_x m_x + (H(u_x))m_{xx}] \, dx \\ &= \int_0^1 H''(u_x)mu_{xx}^2 \, dx. \end{aligned} \tag{6-5}$$

Furthermore, we have

$$- \int_0^1 m^\alpha m_{xx} \, dx = \int_0^1 \alpha m^{\alpha-1} m_x^2 \, dx \tag{6-6}$$

and

$$\int_0^1 -Vm_{xx} \, dx = - \int_0^1 V_{xx}m \, dx \leq \int_0^1 |V_{xx}|m \, dx \leq C \int_0^1 m \, dx \leq C, \tag{6-7}$$

where we used Corollary 5.2.

Finally,

$$\epsilon \int_0^1 (-mm_{xx} + m_{xx}^2 - uu_{xx} + u_{xx}^2) dx = \epsilon \int_0^1 (m_x^2 + m_{xx}^2 + u_x^2 + u_{xx}^2) dx. \tag{6-8}$$

Using (6-3)–(6-8), we get

$$\begin{aligned} \int_0^1 H''(u_x)mu_{xx}^2 dx + \int_0^1 \alpha m^{\alpha-1}m_x^2 dx \\ + \epsilon \int_0^1 (m_x^2 + m_{xx}^2 + u_x^2 + u_{xx}^2) dx = - \int_0^1 Vm_{xx} \leq C. \end{aligned}$$

This completes the proof of (6-1). To conclude the proof of Proposition 6.1, we observe that Assumption 4 implies that  $H''$  is a nonnegative function, which together with (6-1) gives (6-2).  $\square$

### 7. Hölder continuity

We recall that Morrey’s theorem in one dimension [Evans 1998] gives the following result.

**Proposition 7.1.** *Let  $f \in C^1(\mathbb{T})$ . Then,*

$$|f(x) - f(y)| \leq \|f_x\|_{L^2}|x - y|^{1/2} \quad \forall x, y \in \mathbb{T}.$$

**Proposition 7.2.** *Suppose that Assumptions 1–4 hold. Let  $(u, m)$  solve Problem 1. Then,  $u, u_x, m,$  and  $m_x$  are  $\frac{1}{2}$ -Hölder continuous functions with  $L^\infty$ -norms and Hölder constants bounded by  $C/\sqrt{\epsilon}$ , where  $C$  is a universal constant depending only on the constants in Assumptions 1 and 2 and on  $\|V\|_{C^2}$ .*

*Proof.* By Proposition 5.1, we have that

$$\epsilon \int_0^1 (m^2 + u^2 + m_x^2 + u_x^2) dx \leq C, \tag{7-1}$$

where  $C$  is a universal constant depending only on the constants in Assumptions 1 and 2 and on  $\|V\|_{L^\infty}$ .

According to Proposition 7.1, we have

$$|u(x) - u(y)| \leq \|u_x\|_{L^2}|x - y|^{1/2} \quad \forall x, y \in \mathbb{T}. \tag{7-2}$$

Moreover, combining the bound on  $\|u\|_{L^2}$  given by (7-1), the mean-value theorem for definite integrals, and the Hölder continuity given by (7-2), we get the  $L^\infty$  bound on  $u$ . A similar inequality holds for  $m$ . Next, we observe that Proposition 6.1 (see (6-2)) gives bounds for  $\|u_{xx}\|_{L^2}$  and  $\|m_{xx}\|_{L^2}$  of the same type as (7-1). Accordingly, the functions  $u_x$  and  $m_x$  are also  $\frac{1}{2}$ -Hölder continuous, and their  $L^\infty$  norms are

bounded by  $C/\sqrt{\epsilon}$ , where  $C$  depends only on the constants in Assumptions 1 and 2 and on  $\|V\|_{C^2}$ .  $\square$

**Remark 7.3.** Consider [Problem 1](#) with  $V$  replaced by  $\lambda V$  for some  $\lambda \in [0, 1]$ . By revisiting the proofs of [Propositions 5.1](#) and [6.1](#), we can readily check that the bounds stated in these propositions are uniform with respect to  $\lambda \in [0, 1]$ . More precisely, [\(5-1\)](#), [\(6-1\)](#), and [\(6-2\)](#) are still valid for a universal positive constant  $C$  that depends only on the constants in Assumptions 1 and 2 and on  $\|V\|_{C^2}$ . In particular, [Proposition 7.2](#) remains unchanged.

## 8. Higher regularity

The bounds in the previous section give Hölder regularity for any solution  $(u, m)$  of [Problem 1](#) and for its derivatives  $(u_x, m_x)$ . Here, we use [\(2-1\)](#) to improve this result and prove Hölder regularity for  $u_{xx}$  and  $m_{xx}$ .

**Proposition 8.1.** *Suppose that Assumptions 1–5 hold. Let  $(u, m)$  solve [Problem 1](#). Then  $(u, m) \in C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T})$ .*

*Proof.* Solving for  $m - m_{xx}$  in the second equation of [\(2-1\)](#) and replacing the resulting expression in the first equation yields

$$[1 + \epsilon^2 + \epsilon H''(u_x)m]u_{xx} = (1 + \epsilon^2)u + H(u_x) - \epsilon + V(x) - m^\alpha - \epsilon H'(u_x)m_x. \quad (8-1)$$

Because  $H$  is convex, we have  $H''(u_x) \geq 0$ . Consequently,  $1 + \epsilon^2 + \epsilon H''(u_x)m \geq 1 > 0$ . This allows us to rewrite [\(8-1\)](#) as

$$u_{xx} = \frac{(1 + \epsilon^2)u + H(u_x) - \epsilon + V(x) - m^\alpha - \epsilon H'(u_x)m_x}{1 + \epsilon^2 + \epsilon H''(u_x)m}. \quad (8-2)$$

Because  $u, m, u_x,$  and  $m_x$  are  $\frac{1}{2}$ -Hölder continuous and because  $H$  and  $H'$  are locally Lipschitz functions, it follows that

$$(1 + \epsilon^2)u + H(u_x) - \epsilon + V(x) - m^\alpha - \epsilon H'(u_x)m_x$$

is also  $\frac{1}{2}$ -Hölder continuous. Similarly, due to [Assumption 5](#),  $1 + \epsilon^2 + \epsilon H''(u_x)m$  is also  $\frac{1}{2}$ -Hölder continuous and bounded from below by 1. Therefore,  $u_{xx}$  is  $\frac{1}{2}$ -Hölder continuous; thus,  $u \in C^{2,1/2}(\mathbb{T})$ .

Finally, we observe that the second equation in [\(2-1\)](#) is equivalent to

$$m_{xx} = m + \epsilon(u - u_{xx}) - 1 - H''(u_x)mu_{xx} - H'(u_x)m_x. \quad (8-3)$$

Hence, analogous arguments to those used above yield that  $m_{xx}$  is also  $\frac{1}{2}$ -Hölder continuous. Thus,  $m \in C^{2,1/2}(\mathbb{T})$ .  $\square$

### 9. Lower bounds on $m$

Here, we establish our last a priori estimate, which gives lower bounds on  $m$ . We begin by proving an auxiliary result.

**Lemma 9.1.** *Suppose that Assumptions 1–4, 6, and 7 hold. Let  $(u, m)$  solve Problem 1. Then,  $\|\epsilon(u - u_{xx})\|_\infty \leq C\epsilon^{1-\gamma/2}$ , where  $C$  is a universal positive constant depending only on the constants in Assumptions 1, 2, and 7 and on  $\|V\|_{C^2}$ .*

*Proof.* To simplify the notation,  $C$  represents a positive constant depending only on the constants in Assumptions 1, 2, and 7 and on  $\|V\|_{C^2}$  and whose value may change from one instance to another.

Note that  $\max\{\epsilon^{1/2}, \epsilon, \epsilon^{2-\gamma/2}, \epsilon^{3/2}, \epsilon^2\} \leq \epsilon^{1-\gamma/2}$  because  $0 < 1 - \frac{1}{2}\gamma < \frac{1}{2}$  (see Assumption 6).

By Proposition 7.2, we have that  $\|u\|_\infty \leq C/\sqrt{\epsilon}$ . Thus,

$$\|\epsilon u\|_\infty \leq C\epsilon^{1-\gamma/2}. \tag{9-1}$$

Next, we examine  $\|\epsilon u_{xx}\|_\infty$ . The identity (8-2) and the condition  $1 + \epsilon^2 + \epsilon H''(u_x)m > 1$  give

$$\begin{aligned} \|\epsilon u_{xx}\|_\infty &\leq \|\epsilon(1 + \epsilon^2)u\|_\infty + \|\epsilon H(u_x)\|_\infty \\ &\quad + \epsilon^2 + \|\epsilon V\|_\infty + \|\epsilon m^\alpha\|_\infty + \|\epsilon^2 H'(u_x)m_x\|_\infty. \end{aligned} \tag{9-2}$$

By (9-1) and by the boundedness of  $V$ , it follows that

$$\|\epsilon(1 + \epsilon^2)u\|_\infty + \epsilon^2 + \|\epsilon V\|_\infty \leq C\epsilon^{1-\gamma/2}.$$

According to Propositions 5.1 and 6.1, we have

$$\int_0^1 m^{\alpha+1} dx \leq C \quad \text{and} \quad \int_0^1 \alpha m^{\alpha-1} m_x^2 dx = \frac{4\alpha}{(\alpha + 1)^2} \int_0^1 (m^{(\alpha+1)/2})_x^2 dx \leq C.$$

The first integral guarantees that there exists  $x_0 \in \mathbb{T}$  such that  $m^{(\alpha+1)/2}(x_0) \leq C$ . Then, because  $m > 0$  and because  $m \in C^1(\mathbb{T})$ , the second integral together with Proposition 7.1 implies that for all  $x \in \mathbb{T}$ ,

$$0 < m^\alpha(x) = (m^{\alpha/2}(x))^2 \leq (m^{(\alpha+1)/2}(x) - m^{(\alpha+1)/2}(x_0) + m^{(\alpha+1)/2}(x_0) + 1)^2 \leq C.$$

Hence,  $\|\epsilon m^\alpha\|_\infty \leq C\epsilon^{1-\gamma/2}$ .

Assumption 1 and Proposition 7.2 give

$$|H(u_x)| \leq C(1 + \epsilon^{-\gamma/2}).$$

This implies that  $\|\epsilon H(u_x)\|_\infty \leq C\epsilon^{1-\gamma/2}$ .

Combining Assumption 7 with Proposition 7.2 gives the bound

$$|H'(u_x)| \leq C(1 + \epsilon^{-(\gamma-1)/2}).$$

By Proposition 7.2, we have that  $|m_x| \leq C/\sqrt{\epsilon}$ . Therefore,

$$\|\epsilon^2 H'(u_x)m_x\|_\infty \leq C\epsilon^{1-\gamma/2}.$$

Collecting all the estimates proved above, we conclude from (9-1) and (9-2) that

$$\|\epsilon(u - u_{xx})\|_\infty \leq C\epsilon^{1-\gamma/2}. \quad \square$$

**Proposition 9.2.** *Suppose that Assumptions 1–4, 6, and 7 hold. Let*

$$\bar{\epsilon}_0 := \left(\frac{1}{2C}\right)^{2/(2-\gamma)},$$

where  $C$  is the constant given by Lemma 9.1. Let  $(u, m)$  solve Problem 1 with  $0 < \epsilon < \min\{1, \bar{\epsilon}_0\}$ . Then, there exists  $\bar{m} > 0$  such that  $m > \bar{m}$  on  $\mathbb{T}$ . Moreover,  $\bar{m}$  is a universal constant depending only on the constants in Assumptions 1, 2, and 7, on  $\|V\|_{C^2}$ , and on  $\epsilon$ .

*Proof.* Multiplying the second equation in (2-1) by  $1/m$  and integrating with respect to  $x$  in  $[0, 1]$ , we obtain

$$\int_0^1 \left(1 - \frac{m_{xx}}{m} - \frac{(H'(u_x)m)_x}{m}\right) dx = \int_0^1 \left(\frac{1}{m} - \epsilon \frac{u - u_{xx}}{m}\right) dx. \quad (9-3)$$

Integration by parts and periodicity yield

$$\int_0^1 \frac{m_{xx}}{m} dx = \int_0^1 \frac{m_x^2}{m^2} dx.$$

Then, (9-3) can be rewritten as

$$\int_0^1 \left(\frac{1}{m} + \frac{m_x^2}{m^2}\right) dx = 1 + \int_0^1 \frac{\epsilon(u - u_{xx})}{m} dx - \int_0^1 \frac{(H'(u_x)m)_x}{m} dx.$$

Next, we estimate the right-hand side of this identity. By Lemma 9.1, for  $0 < \epsilon < \bar{\epsilon}_0$ , we have  $\|\epsilon(u - u_{xx})\|_\infty < \frac{1}{2}$ . Consequently,

$$\int_0^1 \left(\frac{1}{2m} + \frac{m_x^2}{m^2}\right) dx \leq 1 + \left| \int_0^1 \frac{(H'(u_x)m)_x}{m} dx \right| = 1 + \left| \int_0^1 H'(u_x) \frac{m_x}{m} dx \right|, \quad (9-4)$$

where in the last equality we used the integration by parts formula and the periodicity of  $u_x$ . In view of Cauchy’s inequality, we conclude that

$$\left| \int_0^1 H'(u_x) \frac{m_x}{m} dx \right| \leq \int_0^1 \left| H'(u_x) \frac{m_x}{m} \right| dx \leq \int_0^1 \left( \frac{(H'(u_x))^2}{2} + \frac{m_x^2}{2m^2} \right) dx. \quad (9-5)$$

Invoking Assumptions 6 and 7, we obtain the estimates

$$(H'(u_x))^2 \leq \bar{C}^2(1 + |u_x|^{\gamma-1})^2 \leq 2\bar{C}^2(1 + |u_x|^{2(\gamma-1)}) \leq 2\bar{C}^2(2 + |u_x|^2)$$

in  $\mathbb{T}$ . These estimates, (9-4), (9-5), and Proposition 5.1 yield

$$\int_0^1 \left( \frac{1}{2m} + \frac{m_x^2}{2m^2} \right) dx \leq 1 + \bar{C}^2(2 + C/\epsilon).$$

Consequently, for  $\tilde{C} = 2 + 2\bar{C}^2(2 + C/\epsilon)$ , we obtain the bounds

$$\int_0^1 \frac{1}{m} dx \leq \tilde{C} \quad \text{and} \quad \int_0^1 \frac{m_x^2}{m^2} dx = \int_0^1 (\ln(m))_x^2 dx \leq \tilde{C}.$$

The first bound implies that there exists  $x_0 \in \mathbb{T}$  such that  $1/(m(x_0)) \leq \tilde{C} + 1$ ; that is,  $\ln(m(x_0)) \geq -\ln(\tilde{C} + 1)$ . The second bound, together with Proposition 7.1, implies that for all  $x \in \mathbb{T}$ , the value of  $|\ln(m(x)) - \ln(m(x_0))| \leq \sqrt{\tilde{C}}$ . Hence, for all  $x \in \mathbb{T}$ ,

$$m(x) \geq e^{-\sqrt{\tilde{C}} - \ln(\tilde{C} + 1)}. \quad \square$$

**Remark 9.3.** As in Remark 7.3, the statement of Proposition 9.2 remains unchanged if we replace  $V$  by  $\lambda V$  for some  $\lambda \in [0, 1]$  in Problem 1.

### 10. The linearized operator

Consider the functional,  $F$ , defined for  $(u, m, \lambda) \in C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T}; ]0, \infty[) \times [0, 1]$  by

$$F(u, m, \lambda) = \left[ \begin{array}{l} u - u_{xx} + H(u_x) + \lambda V - m^\alpha - \epsilon(m - m_{xx}) \\ m - m_{xx} - (H'(u_x)m)_x - 1 + \epsilon(u - u_{xx}) \end{array} \right]. \quad (10-1)$$

Note that under Assumption 5, the functional  $F$  is a  $C^1$  map between  $C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T}; ]0, \infty[) \times [0, 1]$  and  $C^{0,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T})$ .

To prove Theorem 2.1, we use the continuation method and show that for every  $\lambda \in [0, 1]$ , the equation

$$F(u, m, \lambda) = 0 \quad (10-2)$$

has a solution,  $(u, m) \in C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T}; ]0, \infty[)$ . Theorem 2.1 then follows by taking  $\lambda = 1$  and by observing that system (2-1) is equivalent to  $F(u, m, 1) = 0$ .

The implicit function theorem plays a crucial role in proving the solvability of (10-2). To use this theorem, for each  $\lambda \in [0, 1]$ , we introduce the linearized operator  $L$  of  $F(\cdot, \cdot, \lambda)$  at  $(u, m) \in C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T}; ]0, \infty[)$ ; that is,

$$\begin{aligned} L(f, v) &= \frac{\partial F}{\partial \mu}(u + \mu v, m + \mu f, \lambda) \Big|_{\mu=0} \\ &= \left[ \begin{array}{l} v - v_{xx} + H'(u_x)v_x - \alpha m^{\alpha-1} f - \epsilon(f - f_{xx}) \\ f - f_{xx} - (H''(u_x)v_x m + H'(u_x)f)_x + \epsilon(v - v_{xx}) \end{array} \right] \end{aligned} \quad (10-3)$$

for  $(f, v) \in C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T})$ . Under Assumption 5 and because  $(u, m) \in C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T}; ]0, \infty[)$ , the operator  $L$  defines a map from  $C^{2,1/2}(\mathbb{T}) \times$

$C^{2,1/2}(\mathbb{T})$  into  $C^{0,1/2}(\mathbb{T}) \times C^{0,1/2}(\mathbb{T})$ . Moreover, this map is continuous and linear. Next, we show that it is also an isomorphism between  $C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T})$  and  $C^{0,1/2}(\mathbb{T}) \times C^{0,1/2}(\mathbb{T})$ .

**Proposition 10.1.** *Suppose that Assumptions 4 and 5 hold. Fix  $\lambda \in [0, 1]$  and assume that  $(u, m) \in C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T}; ]0, \infty[)$  satisfies  $F(u, m, \lambda) = 0$ . Then, the operator  $L$  given by (10-3) is an isomorphism between  $C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T})$  and  $C^{0,1/2}(\mathbb{T}) \times C^{0,1/2}(\mathbb{T})$ .*

*Proof.* To prove the proposition, we begin by applying the Lax–Milgram theorem in  $H^1(\mathbb{T}) \times H^1(\mathbb{T})$ , after which we bootstrap additional regularity. Here, we endow  $H^1(\mathbb{T}) \times H^1(\mathbb{T})$  with the inner product

$$\langle (\theta_1, \theta_2), (\bar{\theta}_1, \bar{\theta}_2) \rangle_{H^1(\mathbb{T}) \times H^1(\mathbb{T})} = \int_0^1 (\theta_1 \bar{\theta}_1 + \theta_2 \bar{\theta}_2 + \theta_{1x} \bar{\theta}_{1x} + \theta_{2x} \bar{\theta}_{2x}) \, dx$$

for  $(\theta_1, \theta_2), (\bar{\theta}_1, \bar{\theta}_2) \in H^1(\mathbb{T}) \times H^1(\mathbb{T})$ .

Consider the bilinear form  $B : (H^1(\mathbb{T}) \times H^1(\mathbb{T})) \times (H^1(\mathbb{T}) \times H^1(\mathbb{T})) \rightarrow \mathbb{R}$  defined for  $(v, f), (w_1, w_2) \in H^1(\mathbb{T}) \times H^1(\mathbb{T})$  by

$$\begin{aligned} B\left(\begin{pmatrix} v \\ f \end{pmatrix}, \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}\right) &= \int_0^1 (f + \epsilon v) w_1 \, dx + \int_0^1 [f_x + H''(u_x) v_x m + H'(u_x) f + \epsilon v_x] w_{1x} \, dx \\ &\quad - \int_0^1 [v + H'(u_x) v_x - \alpha m^{\alpha-1} f - \epsilon f] w_2 \, dx + \int_0^1 (\epsilon f_x - v_x) w_{2x} \, dx. \end{aligned}$$

Note that if  $(v, f) \in C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T})$ , then

$$B\left(\begin{pmatrix} v \\ f \end{pmatrix}, \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}\right) = \int_0^1 [-L_1(f, v) w_2 + L_2(f, v) w_1] \, dx, \tag{10-4}$$

where  $L_1$  and  $L_2$  are the first and second components of  $L$ , respectively.

Next, we prove that  $B$  is coercive and bounded in  $H^1(\mathbb{T}) \times H^1(\mathbb{T})$ . Fix  $(v, f), (w_1, w_2) \in H^1(\mathbb{T}) \times H^1(\mathbb{T})$ . Using the integration by parts formula and the periodicity of  $v$  and  $f$ , we obtain

$$B\left(\begin{pmatrix} v \\ f \end{pmatrix}, \begin{pmatrix} v \\ f \end{pmatrix}\right) = \int_0^1 [\alpha m^{\alpha-1} f^2 + H''(u_x) v_x^2 m + \epsilon(v^2 + v_x^2 + f^2 + f_x^2)] \, dx.$$

Because  $H'' \geq 0$  by Assumption 4 and because  $m > 0$ , we have

$$B\left(\begin{pmatrix} v \\ f \end{pmatrix}, \begin{pmatrix} v \\ f \end{pmatrix}\right) \geq \epsilon \left\| \begin{pmatrix} v \\ f \end{pmatrix} \right\|_{H^1(\mathbb{T}) \times H^1(\mathbb{T})}^2,$$

which proves the coercivity of  $B$ .



Because  $m, u,$  and  $H$  are  $C^{2,1/2}$ -functions on the compact set  $[0, 1]$ , we have that  $m, u, m_x, u_x, u_{xx}, H, H'(u_x),$  and  $H''(u_x)$  are bounded. Therefore, there exists a positive constant,  $C,$  that depends only on these bounds and for which

$$\left| B\left(\begin{pmatrix} v \\ f \end{pmatrix}, \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}\right) \right| \leq C \left\| \begin{pmatrix} v \\ f \end{pmatrix} \right\|_{H^1(\mathbb{T}) \times H^1(\mathbb{T})} \left\| \begin{pmatrix} w_1 \\ w_2 \end{pmatrix} \right\|_{H^1(\mathbb{T}) \times H^1(\mathbb{T})},$$

where we also used Hölder’s inequality. This proves the boundedness of  $B.$

Finally, we fix  $b = (b_1, b_2) \in C^{0,1/2}(\mathbb{T}) \times C^{0,1/2}(\mathbb{T}),$  and we consider the bounded and linear functional  $G : H^1(\mathbb{T}) \times H^1(\mathbb{T}) \rightarrow \mathbb{R}$  defined for  $(w_1, w_2) \in H^1(\mathbb{T}) \times H^1(\mathbb{T})$  by

$$G\left(\begin{pmatrix} w_1 \\ w_2 \end{pmatrix}\right) = \int_0^1 (-b_1 w_2 + b_2 w_1) dx.$$

By the Lax–Milgram theorem, there exists a unique  $(v, f) \in H^1(\mathbb{T}) \times H^1(\mathbb{T})$  such that for all  $(w_1, w_2) \in H^1(\mathbb{T}) \times H^1(\mathbb{T}),$  we have

$$B\left(\begin{pmatrix} v \\ f \end{pmatrix}, \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}\right) = G\left(\begin{pmatrix} w_1 \\ w_2 \end{pmatrix}\right).$$

This is equivalent to saying that for all  $(w_1, w_2) \in H^1(\mathbb{T}) \times H^1(\mathbb{T}),$

$$B\left(\begin{pmatrix} v \\ f \end{pmatrix}, \begin{pmatrix} -w_2 \\ w_1 \end{pmatrix}\right) = G\left(\begin{pmatrix} -w_2 \\ w_1 \end{pmatrix}\right) = \int_0^1 (-b_1 w_1 - b_2 w_2) dx.$$

From this and (10-4), we conclude that  $L(f, v) = b$  has a unique weak solution  $(f, v) \in H^1(\mathbb{T}) \times H^1(\mathbb{T}).$  Because  $b \in C^{0,1/2}(\mathbb{T}) \times C^{0,1/2}(\mathbb{T})$  is arbitrary,  $L$  is injective. To prove surjectivity, it suffices to check that the weak solution of  $L(f, v) = b$  is in  $C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T}).$  This higher regularity follows from a bootstrap argument.

Fix  $b = (b_1, b_2) \in C^{0,1/2}(\mathbb{T}) \times C^{0,1/2}(\mathbb{T})$  and let  $(f, v) \in H^1(\mathbb{T}) \times H^1(\mathbb{T})$  be the weak solution of  $L(f, v) = b$  given by the Lax–Milgram theorem. Then, we have the following identity in the weak sense:

$$v_{xx} = \frac{g}{1 + \epsilon^2 + \epsilon H''(u_x)m}, \tag{10-5}$$

where

$$g = v(1 + \epsilon^2) + H'(u_x)v_x - \alpha m^{\alpha-1} f - \epsilon v_x (H'(u_x)m)_x - \epsilon (H'(u_x)f)_x - \epsilon b_2 - b_1 \in L^2(\mathbb{T}).$$

We recall that  $1 + \epsilon^2 + \epsilon H''(u_x)m > 1.$  Hence,  $v_{xx} \in L^2(\mathbb{T}),$  and so  $v \in H^2(\mathbb{T}).$  Moreover, because

$$f_{xx} = f - (H''(u_x)v_x m)_x - (H'(u_x)f)_x + \epsilon(v - v_{xx}) - b_2 \tag{10-6}$$

in the weak sense, similar arguments yield  $f_{xx} \in L^2(\mathbb{T})$  and  $f \in H^2(\mathbb{T})$ .

So far,  $(f, v) \in C^{1,1/2}(\mathbb{T}) \times C^{1,1/2}(\mathbb{T})$ . This implies that  $g \in C^{0,1/2}(\mathbb{T})$ . Then, using the fact that  $1 + \epsilon^2 + \epsilon H''(u_x)m$  also belongs to  $C^{0,1/2}(\mathbb{T})$  and is bounded from below by 1, from (10-5) it follows that  $v_{xx} \in C^{0,1/2}(\mathbb{T})$ . Consequently, in view of (10-6),  $f_{xx} \in C^{0,1/2}(\mathbb{T})$ . Hence,  $(f, v) \in C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T})$ . Therefore, the unique solution given by the Lax–Milgram theorem is a strong solution with  $C^{2,1/2}$  regularity. Thus,  $L$  is surjective. Because  $L$  is injective and surjective, it is an isomorphism.  $\square$

## 11. Proof of the main theorem

In this last section, we prove [Theorem 2.1](#). We assume that  $\epsilon > 0$  satisfies  $\epsilon < \min\{1, \epsilon_0, \bar{\epsilon}_0\}$ , where  $\epsilon_0$  and  $\bar{\epsilon}_0$  are given by [Propositions 4.1](#) and [9.2](#), respectively.

Let  $F$  be the functional defined in (10-1). For each  $\lambda \in [0, 1]$ , consider the problem of finding  $(u, m) \in C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T}; ]0, \infty[)$  satisfying (10-2). From [Propositions 4.1](#) and [8.1](#), such a pair  $(u, m)$  exists for  $\lambda = 0$ . Next, using the continuation method, we prove that this is true not only for  $\lambda = 0$  but also for all  $\lambda \in [0, 1]$ .

More precisely, let  $\Lambda$  be the set of values  $\lambda \in [0, 1]$  for which (10-2) has a solution  $(u, m) \in C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T})$  with  $m \geq \bar{m}$  in  $\mathbb{T}$ , where  $\bar{m} > 0$  is given by [Proposition 9.2](#). Note that  $\bar{m}$  does not depend on  $\lambda$  (see [Remark 9.3](#)). As we just argued,  $\Lambda$  is a nonempty set. In the subsequent two propositions, we show that  $\Lambda$  is a closed and open subset of  $[0, 1]$ . Consequently,  $\Lambda = [0, 1]$ .

**Proposition 11.1.** *Suppose that Assumptions 1–7 hold. Then,  $\Lambda$  is a closed subset of  $[0, 1]$ .*

*Proof.* Let  $(\lambda^n)_{n \in \mathbb{N}} \subset \Lambda$  and  $\lambda \in [0, 1]$  be such that  $\lim_{n \rightarrow \infty} \lambda^n = \lambda$ . We claim that  $\lambda \in \Lambda$ .

By definition of  $\Lambda$ , for each  $n \in \mathbb{N}$ , there exists  $(u^n, m^n) \in C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T})$  satisfying (10-2) and  $m^n \geq \bar{m}$  in  $\mathbb{T}$ . Then, by [Proposition 7.2](#) (also see [Remark 7.3](#)),  $(u^n)_{n \in \mathbb{N}}$ ,  $(m^n)_{n \in \mathbb{N}}$ ,  $(u_x^n)_{n \in \mathbb{N}}$ , and  $(m_x^n)_{n \in \mathbb{N}}$  are uniformly bounded in  $C^{0,1/2}(\mathbb{T})$ . Consequently, by the Arzelà–Ascoli theorem, we can find  $(u, m, \tilde{u}, \tilde{m}) \in C^{0,1/2}(\mathbb{T}) \times C^{0,1/2}(\mathbb{T}) \times C^{0,1/2}(\mathbb{T}) \times C^{0,1/2}(\mathbb{T})$  such that, up to a subsequence that we do not relabel,

$$\lim_{n \rightarrow \infty} \|(u^n, m^n, u_x^n, m_x^n) - (u, m, \tilde{u}, \tilde{m})\|_\infty = 0. \quad (11-1)$$

We now recall that if  $(w^n)_{n \in \mathbb{N}}$  is a sequence of differentiable functions on  $[0, 1]$  such that  $(w^n)_{n \in \mathbb{N}}$  converges uniformly to some  $w$  on  $[0, 1]$  and such that  $(w_x^n)_{n \in \mathbb{N}}$  converges uniformly on  $[0, 1]$ , then  $w_x = \lim_{n \rightarrow \infty} w_x^n$  on  $[0, 1]$ . Consequently, by (11-1), we have  $\tilde{u} = u_x$  and  $\tilde{m} = m_x$ .

Next, we show that  $(u_{xx}^n)_{n \in \mathbb{N}}$  and  $(m_{xx}^n)_{n \in \mathbb{N}}$  are also uniformly convergent sequences on  $[0, 1]$ . In view of (8-2), we have for every  $n \in \mathbb{N}$ ,

$$u_{xx}^n = \frac{(1 + \epsilon^2)u^n + H(u_x^n) - \epsilon + \lambda^n V(x) - (m^n)^\alpha - \epsilon H'(u_x^n)m_x^n}{1 + \epsilon^2 + \epsilon H''(u_x^n)m^n}. \tag{11-2}$$

By Assumption 5 and by the uniform convergence of  $(u^n, m^n, \lambda^n, u_x^n, m_x^n)_{n \in \mathbb{N}}$  to  $(u, m, \lambda, u_x, m_x)$  on  $[0, 1]$ , it follows from (11-2) that  $(u_{xx}^n)_{n \in \mathbb{N}}$  converges uniformly on  $[0, 1]$ . Then, the limit of  $(u_{xx}^n)_{n \in \mathbb{N}}$  is necessarily  $u_{xx}$ . Analogous arguments (see (8-3)) give that  $(m_{xx}^n)_{n \in \mathbb{N}}$  converges uniformly to  $m_{xx}$  on  $[0, 1]$ . Thus,  $(u, m) \in C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T}; ]0, \infty[)$ . Moreover,  $\lim_{n \rightarrow \infty} F(u^n, m^n, \lambda^n) = F(u, m, \lambda)$ . Finally, because for all  $n \in \mathbb{N}$ , the functional  $F(u^n, m^n, \lambda^n) = 0$  and  $m^n \geq \bar{m}$  in  $\mathbb{T}$ , we have that  $F(u, m, \lambda) = 0$  and  $m \geq \bar{m}$  in  $\mathbb{T}$ . Hence,  $\lambda \in \Lambda$ .  $\square$

**Proposition 11.2.** *Suppose that Assumptions 1–7 hold. Then,  $\Lambda$  is an open subset of  $[0, 1]$ .*

*Proof.* Let  $\lambda_0 \in \Lambda$ . Then, there exists  $(u_0, m_0) \in C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T})$  satisfying  $F(u_0, m_0, \lambda_0) = 0$  and  $m_0 \geq \bar{m}$  in  $\mathbb{T}$ . By Proposition 10.1 and by the implicit function theorem in Banach spaces (see, for example, [Dieudonné 1960]), we can find  $\delta > 0$  such that, for every  $\lambda^* \in ]\lambda - \lambda_0, \lambda + \lambda_0[$ , there exists  $(u^*, m^*) \in C^{2,1/2}(\mathbb{T}) \times C^{2,1/2}(\mathbb{T})$  satisfying  $F(u^*, m^*, \lambda^*) = 0$  and  $m^* \geq \bar{m}$  in  $\mathbb{T}$ . Moreover, the implicit function theorem also guarantees that the map  $\lambda^* \mapsto m^*$  is continuous. Hence, if  $\delta$  is small enough, we have  $m^* > 0$  in  $\mathbb{T}$ . Then, Proposition 9.2 gives  $m^* > \bar{m}$  in  $\mathbb{T}$ . Therefore,  $\lambda^* \in \Lambda$  and, consequently,  $\Lambda$  is open.  $\square$

Finally, we sum up the proof of our main result.

*Proof of Theorem 2.1.* Let  $\epsilon > 0$  be such that  $\epsilon < \min\{1, \epsilon_0, \bar{\epsilon}_0\}$ , where  $\epsilon_0$  is given by Proposition 4.1 and where  $\bar{\epsilon}_0$  is given by Proposition 9.2.

Propositions 11.1 and 11.2 give that  $\Lambda$  is a relatively open and closed set in  $[0, 1]$ . It is a nonempty set due to Propositions 4.1, 8.1, and 9.2. Hence,  $\Lambda = [0, 1]$ . Finally, we observe that Theorem 2.1 corresponds to the  $\lambda = 1$  case.  $\square$

### Acknowledgements

This summer camp would not have been possible without major help and support from KAUST. David Yeh and his team at the Visiting Student Research Program did a fantastic job in organizing all the logistics. In addition, the CEMSE Division provided valuable additional support. Finally, we would like to thank the faculty, research scientists, postdocs and Ph.D. students who, in the first three weeks of the semester, made time to give courses, lectures, and work with the students.

## References

- [Cardaliaguet et al. 2015] P. Cardaliaguet, P. J. Graber, A. Porretta, and D. Tonon, “Second order mean field games with degenerate diffusion and local coupling”, *NoDEA Nonlinear Differential Equations Appl.* **22**:5 (2015), 1287–1317. [MR](#) [Zbl](#)
- [Cirant 2015] M. Cirant, “Multi-population mean field games systems with Neumann boundary conditions”, *J. Math. Pures Appl.* (9) **103**:5 (2015), 1294–1315. [MR](#) [Zbl](#)
- [Dieudonné 1960] J. Dieudonné, *Foundations of modern analysis*, Pure and applied mathematics **10**, Academic Press, New York, 1960. Reprinted 1969 as v. 1 of *Treatise on analysis*. [MR](#) [Zbl](#)
- [Evans 1998] L. C. Evans, *Partial differential equations*, Graduate Studies in Mathematics **19**, American Mathematical Society, Providence, RI, 1998. [MR](#) [Zbl](#)
- [Ferreira and Gomes 2015] R. Ferreira and D. Gomes, “Existence of weak solutions to stationary mean-field games through variational inequalities”, Preprint, 2015. [arXiv](#)
- [Gomes and Mitake 2015] D. A. Gomes and H. Mitake, “Existence for stationary mean-field games with congestion and quadratic Hamiltonians”, *NoDEA Nonlinear Differential Equations Appl.* **22**:6 (2015), 1897–1910. [MR](#) [Zbl](#)
- [Gomes and Patrizi 2015] D. A. Gomes and S. Patrizi, “Obstacle mean-field game problem”, *Interfaces Free Bound.* **17**:1 (2015), 55–68. [MR](#) [Zbl](#)
- [Gomes and Pimentel 2015] D. A. Gomes and E. Pimentel, “Time-dependent mean-field games with logarithmic nonlinearities”, *SIAM J. Math. Anal.* **47**:5 (2015), 3798–3812. [MR](#) [Zbl](#)
- [Gomes and Pimentel 2016] D. A. Gomes and E. Pimentel, “Local regularity for mean-field games in the whole space”, *Minimax Theory Appl.* **1**:1 (2016), 65–82. [MR](#) [Zbl](#)
- [Gomes and Ribeiro 2013] D. Gomes and R. Ribeiro, “Mean field games with logistic population dynamics”, in *52nd IEEE Conference on Decision and Control* (Florence), IEEE, Piscataway, NJ, 2013.
- [Gomes and Sánchez Morgado 2014] D. Gomes and H. Sánchez Morgado, “A stochastic Evans–Aronsson problem”, *Trans. Amer. Math. Soc.* **366**:2 (2014), 903–929. [MR](#) [Zbl](#)
- [Gomes and Saúde 2014] D. A. Gomes and J. Saúde, “Mean field games models — a brief survey”, *Dyn. Games Appl.* **4**:2 (2014), 110–154. [MR](#)
- [Gomes et al. 2012] D. A. Gomes, G. E. Pires, and H. Sánchez-Morgado, “A-priori estimates for stationary mean-field games”, *Netw. Heterog. Media* **7**:2 (2012), 303–314. [MR](#) [Zbl](#)
- [Gomes et al. 2014] D. A. Gomes, S. Patrizi, and V. Voskanyan, “On the existence of classical solutions for stationary extended mean field games”, *Nonlinear Anal.* **99** (2014), 49–79. [MR](#) [Zbl](#)
- [Gomes et al. 2015] D. A. Gomes, E. A. Pimentel, and H. Sánchez-Morgado, “Time-dependent mean-field games in the subquadratic case”, *Comm. Partial Differential Equations* **40**:1 (2015), 40–76. [MR](#) [Zbl](#)
- [Gomes et al. 2016] D. A. Gomes, E. Pimentel, and H. Sánchez-Morgado, “Time-dependent mean-field games in the superquadratic case”, *ESAIM Control Optim. Calc. Var.* **22**:2 (2016), 562–580. [MR](#) [Zbl](#)
- [Graber 2015] J. Graber, “Weak solutions for mean field games with congestion”, Preprint, 2015. [arXiv](#)
- [Huang et al. 2006] M. Huang, R. P. Malhamé, and P. E. Caines, “Large population stochastic dynamic games: closed-loop McKean–Vlasov systems and the Nash certainty equivalence principle”, *Commun. Inf. Syst.* **6**:3 (2006), 221–251. [MR](#) [Zbl](#)

- [Huang et al. 2007] M. Huang, P. E. Caines, and R. P. Malhamé, “Large-population cost-coupled LQG problems with nonuniform agents: individual-mass behavior and decentralized  $\epsilon$ -Nash equilibria”, *IEEE Trans. Automat. Control* **52**:9 (2007), 1560–1571. [MR](#)
- [Lasry and Lions 2006a] J.-M. Lasry and P.-L. Lions, “Jeux à champ moyen, I: le cas stationnaire”, *C. R. Math. Acad. Sci. Paris* **343**:9 (2006), 619–625. [MR](#) [Zbl](#)
- [Lasry and Lions 2006b] J.-M. Lasry and P.-L. Lions, “Jeux à champ moyen, II: horizon fini et contrôle optimal”, *C. R. Math. Acad. Sci. Paris* **343**:10 (2006), 679–684. [MR](#) [Zbl](#)
- [Lasry and Lions 2007] J.-M. Lasry and P.-L. Lions, “Mean field games”, *Jpn. J. Math.* **2**:1 (2007), 229–260. [MR](#) [Zbl](#)
- [Lions 2012] P. Lions, “Course on mean-field games”, 2012, <http://www.ima.umn.edu/2012-2013/sw11.12-13.12/>.
- [Pimentel and Voskanyan 2015] E. Pimentel and V. Voskanyan, “Regularity for second-order stationary mean-field games”, Preprint, 2015. To appear in *Indiana Univ. Math. J.* [arXiv](#)
- [Porretta 2014] A. Porretta, “On the planning problem for the mean field games system”, *Dyn. Games Appl.* **4**:2 (2014), 231–256. [MR](#) [Zbl](#)
- [Porretta 2015] A. Porretta, “Weak solutions to Fokker–Planck equations and mean field games”, *Arch. Ration. Mech. Anal.* **216**:1 (2015), 1–62. [MR](#) [Zbl](#)

Received: 2016-01-12

Revised: 2016-03-21

Accepted: 2016-04-15

[nojoud16@gmail.com](mailto:nojoud16@gmail.com)*Effat University, Jeddah 23523, Saudi Arabia*[elena.bachini@studenti.unipd.it](mailto:elena.bachini@studenti.unipd.it)*Department of Mathematics, University of Padova, I-35121 Padova, Italy*[a\\_chapouto@hotmail.com](mailto:a_chapouto@hotmail.com)*University of Coimbra, 5400-000 Chaves, Portugal*[rita.ferreira@kaust.edu.sa](mailto:rita.ferreira@kaust.edu.sa)*CEMSE Division, King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Saudi Arabia*[diogo.gomes@kaust.edu.sa](mailto:diogo.gomes@kaust.edu.sa)*CEMSE Division, King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Saudi Arabia*[daniela.jordao@hotmail.com](mailto:daniela.jordao@hotmail.com)*University of Coimbra, 3001-501 Coimbra, Portugal*[davidsevangalista@gmail.com](mailto:davidsevangalista@gmail.com)*CEMSE Division, King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Saudi Arabia*[bbavo96@yahoo.com](mailto:bbavo96@yahoo.com)*Faculty of Mathematics and Mechanics, Yerevan State University, Republic of Armenia, Yerevan, 0025, 1 Alex Manoogian, 0025 Yerevan, Armenia*[laterio@gmail.com](mailto:laterio@gmail.com)*University of Buenos Aires, 1414CKS Buenos Aires, Argentina*[levonnurbekian@gmail.com](mailto:levonnurbekian@gmail.com)*CEMSE Division, King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Saudi Arabia*[pagliagiorgia37@gmail.com](mailto:pagliagiorgia37@gmail.com)*University of Verona, I-37045 Legnago, Italy*[piccio991@hotmail.it](mailto:piccio991@hotmail.it)*University of Rome Tor Vergata, I-35121 Padova, Italy*[sagar.dipak@gmail.com](mailto:sagar.dipak@gmail.com)*University of Coimbra, 4465-040 S. Mamede de Infesta, Portugal*

- [mariana.prazeres.1@kaust.edu.sa](mailto:mariana.prazeres.1@kaust.edu.sa) *CEMSE Division, King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Saudi Arabia*
- [joao.feliciodosreis@kaust.edu.sa](mailto:joao.feliciodosreis@kaust.edu.sa) *CEMSE Division, King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Saudi Arabia*
- [amr\\_aac@hotmail.com](mailto:amr_aac@hotmail.com) *University of Coimbra, 3040-109 Coimbra, Portugal*
- [orlando\\_andres2005@hotmail.com](mailto:orlando_andres2005@hotmail.com) *University of Lisbon, 9000-690 Funchal, Portugal*
- [maria\\_sargsyan@edu.aua.am](mailto:maria_sargsyan@edu.aua.am) *American University of Armenia, 0025 Yerevan, Armenia*
- [ttoommxx@gmail.com](mailto:ttoommxx@gmail.com) *University of Verona, via Brunelleschi 1, I-37178 Verona, Italy*
- [larrysong@zju.edu.cn](mailto:larrysong@zju.edu.cn) *MIT, Cambridge, MA 02140, United States*
- [ken5terai@akane.waseda.jp](mailto:ken5terai@akane.waseda.jp) *Waseda University, Saitama 338-0001, Japan*
- [ryota-tomisaki@fuji.waseda.jp](mailto:ryota-tomisaki@fuji.waseda.jp) *Waseda University, 210-10 Nase, Totsuka, Yokohama 245-00051, Japan*
- [bigquantum@ciencias.unam.mx](mailto:bigquantum@ciencias.unam.mx) *National Autonomous University of Mexico, Cerro Copor #16, Campestre Churubusco, 02100 Mexico City,, Mexico*
- [vartanvos@gmail.com](mailto:vartanvos@gmail.com) *CEMSE Division, King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Saudi Arabia*
- [xianjin.yang@kaust.edu.sa](mailto:xianjin.yang@kaust.edu.sa) *CEMSE Division, King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Saudi Arabia*

## INVOLVE YOUR STUDENTS IN RESEARCH

*Involve* showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

### MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

### BOARD OF EDITORS

Colin Adams	Williams College, USA	Suzanne Lenhart	University of Tennessee, USA
John V. Baxley	Wake Forest University, NC, USA	Chi-Kwong Li	College of William and Mary, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Robert B. Lund	Clemson University, USA
Martin Bohner	Missouri U of Science and Technology, USA	Gaven J. Martin	Massey University, New Zealand
Nigel Boston	University of Wisconsin, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA	Emil Minchev	Ruse, Bulgaria
Pietro Cerone	La Trobe University, Australia	Frank Morgan	Williams College, USA
Scott Chapman	Sam Houston State University, USA	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Joshua N. Cooper	University of South Carolina, USA	Zuhair Nashed	University of Central Florida, USA
Jem N. Corcoran	University of Colorado, USA	Ken Ono	Emory University, USA
Toka Diagana	Howard University, USA	Timothy E. O'Brien	Loyola University Chicago, USA
Michael Dorff	Brigham Young University, USA	Joseph O'Rourke	Smith College, USA
Sever S. Dragomir	Victoria University, Australia	Yuval Peres	Microsoft Research, USA
Behrouz Emamizadeh	The Petroleum Institute, UAE	Y.-F. S. Pétermann	Université de Genève, Switzerland
Joel Foisy	SUNY Potsdam, USA	Robert J. Plemmons	Wake Forest University, USA
Erin W. Fulp	Wake Forest University, USA	Carl B. Pomerance	Dartmouth College, USA
Joseph Gallian	University of Minnesota Duluth, USA	Vadim Ponomarenko	San Diego State University, USA
Stephan R. Garcia	Pomona College, USA	Bjorn Poonen	UC Berkeley, USA
Anant Godbole	East Tennessee State University, USA	James Propp	U Mass Lowell, USA
Ron Gould	Emory University, USA	József H. Przytycki	George Washington University, USA
Andrew Granville	Université Montréal, Canada	Richard Rebarber	University of Nebraska, USA
Jerrold Griggs	University of South Carolina, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Jim Haglund	University of Pennsylvania, USA	James A. Sellers	Penn State University, USA
Johnny Henderson	Baylor University, USA	Andrew J. Sterge	Honorary Editor
Jim Hoste	Pitzer College, USA	Ann Trenk	Wellesley College, USA
Natalia Hritonenko	Prairie View A&M University, USA	Ravi Vakil	Stanford University, USA
Glenn H. Hurlbert	Arizona State University, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
Charles R. Johnson	College of William and Mary, USA	Ram U. Verma	University of Toledo, USA
K. B. Kulasekera	Clemson University, USA	John C. Wierman	Johns Hopkins University, USA
Gerry Ladas	University of Rhode Island, USA	Michael E. Zieve	University of Michigan, USA

### PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or [msp.org/involve](http://msp.org/involve) for submission instructions. The subscription price for 2017 is US \$175/year for the electronic version, and \$235/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

# involve

2017

vol. 10

no. 3

Dynamics of vertical real rhombic Weierstrass elliptic functions	361
LORELEI KOSS AND KATIE ROY	
Pattern avoidance in double lists	379
CHARLES CRATTY, SAMUEL ERICKSON, FREHIWET NEGASSI AND LARA PUDWELL	
On a randomly accelerated particle	399
MICHELLE NUNO AND JUHI JANG	
Reeb dynamics of the link of the $A_n$ singularity	417
LEONARDO ABBRESCIA, IRIT HUQ-KURUVILLA, JO NELSON AND NAWAZ SULTANI	
The vibration spectrum of two Euler–Bernoulli beams coupled via a dissipative joint	443
CHRIS ABRIOLA, MATTHEW P. COLEMAN, AGLIKA DARAKCHIEVA AND TYLER WALES	
Loxodromes on hypersurfaces of revolution	465
JACOB BLACKWOOD, ADAM DUKEHART AND MOHAMMAD JAVAHERI	
Existence of positive solutions for an approximation of stationary mean-field games	473
NOJOOD ALMAYOUF, ELENA BACHINI, ANDREA CHAPOUTO, RITA FERREIRA, DIOGO GOMES, DANIELA JORDÃO, DAVID EVANGELISTA JUNIOR, AVETIK KARAGULYAN, JUAN MONASTERIO, LEVON NURBEKYAN, GIORGIA PAGLIAR, MARCO PICCIRILLI, SAGAR PRATAPSI, MARIANA PRAZERES, JOÃO REIS, ANDRÉ RODRIGUES, ORLANDO ROMERO, MARIA SARGSYAN, TOMMASO SENECCI, CHULIANG SONG, KENGO TERAJ, RYOTA TOMISAKI, HECTOR VELASCO-PEREZ, VARDAN VOSKANYAN AND XIANJIN YANG	
Discrete dynamics of contractions on graphs	495
OLENA OSTAPYUK AND MARK RONNENBERG	
Tiling annular regions with skew and T-tetrominoes	505
AMANDA BRIGHT, GREGORY J. CLARK, CHARLES DUNN, KYLE EVITTS, MICHAEL P. HITCHMAN, BRIAN KEATING AND BRIAN WHETTER	
A bijective proof of a $q$ -analogue of the sum of cubes using overpartitions	523
JACOB FORSTER, KRISTINA GARRETT, LUKE JACOBSEN AND ADAM WOOD	
Ulrich partitions for two-step flag varieties	531
IZZET COSKUN AND LUKE JASKOWIAK	